



Button at top left
quickly takes
user to main menu



Scrolling alphabet indexer
on the right. Tap a letter
on the right,
and list on left scrolls to
that letter (like iPhone
Contacts)





Contact info and
EULA could go under
About Us

Swipe the partially
hidden screen
to return to it (or
tap it on menu)

2 new
notifications

If we add the Bookmarks and Search options, they will probably go to the top





Condition Quick Pick



Look up a condition:

B

Bacterial Vaginosis

C

Candidiasis, Vulvovaginal

Cervicitis

Chancroid

Chlamydia

E


Epididymitis

G


Genital Warts

A
B
C
D
E
F
G
H
I
J
K
L
M
N
O
P
Q
R
S
T
U
V
W
X
Y
Z

User selects Chlamydia





If applicable, a notification appears in a banner at the bottom (only one line of text, then "..."). Tap it to read entire notification.



Once user taps the notification banner, it displays all the text. Once user taps it again or anywhere in the white space, it collapses to an icon. See next screen...

Persistent, non-obtrusive icon stays here. It is translucent, so can see content beneath. Tapping this makes the notification appear on the screen again.





In the event that a screen has multiple notifications, the user can navigate through them. (This design assumes that multiple notifications are not the common case, and if there are multiple notifications, there are only a few (2 or 3); however, if this is untrue, we will need to rethink this design).

ABC 3G

12:17 PM



Condition Quick Pick



Chlamydia

[See related content ...](#)

Adolescents & Adults

Recommended Regimens

(text)

Alternative Regimens

(text)




Treatments



Description




Photos



User taps
'See related
content...'

This is a new page. Tapping 'Done' takes user right back to where he/she was



ABC 3G

12:17 PM

Done

Related Content



Chlamydia

Partner Management

When patients diagnosed with chlamydia or gonorrhea indicate that their partners are unlikely to seek evaluation and treatment, providers can offer patient-delivered partner therapy (PDPT), a form of expedited partner therapy (EPT) in which partners of infected persons are treated without previous medical evaluation or prevention counseling. Because EPT might be prohibited in some states and is the topic of ongoing legislation in others (67), providers should visit www.cdc.gov/std/ept to obtain updated information for their individual jurisdiction.

ABC 3G

12:17 PM



Condition Quick Pick



Chlamydia

[See related content ...](#)

Adolescents & Adults >

Among Infants >

Among Children >



Treatments



Description



Photos

ABC 3G

12:17 PM



Condition Quick Pick



Chlamydia

[See related content ...](#)

Adolescents & Adults

Diagnostic Considerations



Treatment



Follow-Up



Management of Sex Partners



Special Considerations



Treatments




Description



Photos

Example of persistent icon w/ transparency so text beneath can be seen.

If needed, would it be possible for the user to touch anywhere on the screen (except a control), and the icon disappears until user touches non-control area on screen again? And the icon could be reset to appear once user leaves this tagged topic?



If user had picked
Special Considerations,
which has sub-topics





ABC 3G

12:17 PM



Condition Quick Pick



Chlamydia

(Chlamydia photo(s) here)




Treatments



Description



Photos



Look

B

Bact

C

Cand

Cerv

Chan

Chla

E

Epidi

G

Geni


User taps
Notifications




'New' notifications are the ones indicated in bold w/ an icon. The user hasn't read it yet.

A user can tap a notification to read the entire notification. From there, user can opt to view the corresponding screen where it appears. (see next screen)


User cannot permanently delete a notification from here. Possibly could in settings???



If user taps a notification (on the previous screen), he/she will come to a screen like this. The entire notification is displayed, as well as the topics to which it applies. The user can select a topic, and see it on the corresponding screen. Topics that have not yet been accessed are in bold (e.g., **Gonorrhea** and **HPV** here).



Example of notification that applies to a single topic.



ABC 3G

12:17 PM



Full STD Guidelines



Summary



Introduction



Methods



Clinical Prevention Guidance



Special Populations



HIV Infection: Detection,
Counseling, and Referral



Diseases Characterized by Genital,
Anal, or Perianal Ulcers



Management of Persons Who Have
a History of Penicillin Allergy



Diseases Characterized by
Urethritis and Cervicitis



Chlamydial Infections




Gonococcal Infections



Diseases Characterized by Vaginal





ABC 3G

12:17 PM



Full STD Guidelines



Clinical Prevention Guidance

Overview



STD/HIV Prevention Counseling



Prevention Methods




Partner Management



Reporting and Confidentiality





ABC 3G

12:17 PM



Full STD Guidelines



Prevention Methods

- Abstinence & Reduction of number of Sex Partners >
- Pre-exposure Vaccination >
- Male Condoms >
- Female Condoms >
- Cervical Diaphragms >
- Topical Microbicides & Spermicides >
- Condoms & n-9 Vaginal Spermicides >
- Rectal Use of n-9 Spermicides >
- Nonbarrier Contraception, Surgical Sterilization, and Hysterectomy >
- Male Circumcision >
- Emergency Contraception (EC) >

This is getting more granular than the Guidelines TOC

ABC 3G

12:17 PM



Full STD Guidelines



Pre-exposure Vaccination

Pre-exposure vaccination is one of the most effective methods for preventing transmission of some STDs. Two human papillomavirus (HPV) vaccines are available for females aged 9-26 years to prevent cervical precancer and cancer (15,16): the quadrivalent HPV vaccine (Gardasil) and the bivalent HPV vaccine (Cervarix). Gardasil also prevents genital warts. Routine vaccination of females aged 11 or 12 years is recommended with either vaccine, as is catch-up vaccination for females aged 13-26 years. Gardasil can be administered to males aged 9-26 years to prevent genital warts (17). Details regarding HPV vaccination are available at



Notification: Pre-exposure vaccines...

ABC 3G

12:17 PM



Full STD Guidelines



Pre-exposure Vaccination

Pre-exposure vaccination is one of the most effective methods for preventing transmission of some STDs. Two human papillomavirus (HPV) vaccines are available for females aged 9-26 years to prevent cervical precancer and cancer (15,16): the quadrivalent HPV vaccine (Gardasil) and the bivalent HPV vaccine (Cervarix). Gardasil also prevents genital warts. Routine vaccination of females aged 11 or 12 years is recommended with either vaccine, as is catch-up vaccination for females aged 13-26 years. Gardasil can be administered to males aged 9-26 years to



Notification: Pre-exposure vaccines orem ipsum dolor sit amet, consectetur quid pro quo ipsum lorem.

ABC 3G

12:17 PM



Full STD Guidelines




Pre-exposure Vaccination

Pre-exposure vaccination is one of the most effective methods for preventing transmission of some STDs. Two human papillomavirus (HPV) vaccines are available for females aged 9-26 years to prevent cervical precancer and cancer (15,16): the quadrivalent HPV vaccine (Gardasil) and the bivalent HPV vaccine (Cervarix). Gardasil also prevents genital warts. Routine vaccination of females aged 11 or 12 years is recommended with either vaccine, as is catch-up vaccination for females aged 13-26 years. Gardasil can be administered to males aged 9-26 years to prevent genital warts (17). Details regarding HPV vaccination are available at



www.cdc.gov/std/hpv. Hepatitis B



Launch the full PDF
(access this from
Main Menu).
PDF Metadata
(size, last updated,
etc.) could go here
too.

