



Button at top left  
quickly takes  
user to main menu




Scrolling alphabet indexer  
on the right. Tap a letter  
on the right,  
and list on left scrolls to  
that letter (like iPhone  
Contacts)

For Iteration 1,  
only the top 3  
options will work



Swipe the partially  
hidden screen  
to return to it (or  
tap it on menu)

User navigates to  
Full STD Guidelines  
from main menu



ABC 3G

02:32 PM



## Full STD Guidelines

Summary	>
Introduction	>
Methods	>
Clinical Prevention Guidance	>
Special Populations	>
HIV Infection: Detection, Counseling, and Referral	>
Diseases Characterized by Genital, Anal, or Perianal Ulcers	>
Management of Persons Who Have a History of Penicillin Allergy	>
Diseases Characterized by Urethritis and Cervicitis	>
Chlamydial Infections	>
Gonococcal Infections	>
Diseases Characterized by Vaginal	>

ABC 3G

02:32 PM



## Full STD Guidelines



### Clinical Prevention Guidance

Overview



STD/HIV Prevention Counseling



Prevention Methods




Partner Management



Reporting and Confidentiality





ABC 3G

02:32 PM



## Full STD Guidelines



### Prevention Methods

- Abstinence & Reduction of number of Sex Partners >
- Pre-exposure Vaccination >
- Male Condoms >
- Female Condoms >
- Cervical Diaphragms >
- Topical Microbicides & Spermicides >
- Condoms & n-9 Vaginal Spermicides >
- Rectal Use of n-9 Spermicides >
- Nonbarrier Contraception, Surgical Sterilization, and Hysterectomy >
- Male Circumcision >
- Emergency Contraception (EC) >

This is getting more granular than the Guidelines TOC

ABC 3G

02:32 PM




## Full STD Guidelines



### Pre-exposure Vaccination


Pre-exposure vaccination is one of the most effective methods for preventing transmission of some STDs. Two human papillomavirus (HPV) vaccines are available for females aged 9-26 years to prevent cervical precancer and cancer (15,16): the quadrivalent HPV vaccine (Gardasil) and the bivalent HPV vaccine (Cervarix). Gardasil also prevents genital warts. Routine vaccination of females aged 11 or 12 years is recommended with either vaccine, as is catch-up vaccination for females aged 13-26 years. Gardasil can be administered to males aged 9-26 years to prevent genital warts (17). Details regarding HPV vaccination are available at [www.cdc.gov/std/hpv](http://www.cdc.gov/std/hpv). Hepatitis B

User navigates to  
PDF of STD  
Guidelines from  
main menu






Launch the full PDF  
(access this from  
Main Menu).  
PDF Metadata  
(size, last updated,  
etc.) could go here  
too.



**Are any of the following wireframes in Iteration 1? ...**

(these all reflect functionality for the Condition Quick Pick)



ABC 3G

02:32 PM



## Condition Quick Pick

# Chlamydia

Adolescents & Adults



Among Infants



Among Children



Treatments



Description

ABC 3G

02:32 PM



## Condition Quick Pick

# Chlamydia

Adolescents & Adults



Among Infants



Among Children



Treatments



Description

ABC 3G

02:32 PM



## Condition Quick Pick



# Chlamydia

### Adolescents & Adults

#### Recommended Regimens

(text)

#### Alternative Regimens

(text)



Treatments



Description

ABC 3G

02:32 PM



## Condition Quick Pick

# Chlamydia

Adolescents & Adults >

Among Infants >

Among Children >



Treatments



Description

ABC 3G

02:32 PM



## Condition Quick Pick



# Chlamydia

Adolescents & Adults

Diagnostic Considerations

Treatment



Follow-Up



Management of Sex Partners



Special Considerations



Treatments



Description



ABC 3G

02:32 PM



## Condition Quick Pick



# Chlamydia

### Diagnostic Considerations

*C. trachomatis* urogenital infection in women can be diagnosed by testing urine or by collecting swab specimens from the endocervix or vagina. Diagnosis of *C. trachomatis* urethral infection in men can be made by testing a urethral swab or urine specimen. Rectal *C. trachomatis* infections in persons that engage in receptive anal intercourse can be diagnosed by testing a rectal swab specimen. NAATs, cell culture, direct immunofluorescence, EIA, and nucleic acid hybridization tests are available for the



Treatments



Description

If user had picked  
Special Considerations,  
which has sub-topics

