



Button at top left quickly takes user to main menu



Scrolling alphabet indexer on the right. Tap a letter on the right, and list on left scrolls to that letter (like iPhone Contacts)





Contact info and  
EULA could go under  
About Us


Swipe the partially  
hidden screen  
to return to it (or  
tap it on menu)

2 new  
notifications


If we add the Bookmarks and Search options, they will probably go to the top








If applicable, a notification appears in a banner at the bottom (only one line of text, then "..."). Tap it to read entire notification.



Once user taps the notification banner, it displays all the text. Once user taps it again or anywhere in the white space, it collapses to an icon. See next screen...



Persistent, non-obtrusive icon stays here. It is translucent, so can see content beneath. Tapping this makes the notification appear on the screen again.

ABC 3G

02:32 PM



## Condition Quick Pick



# Chlamydia

[See related content ...](#)

### Adolescents & Adults

#### Recommended Regimens

(text)

#### Alternative Regimens

(text)



Treatments




Description




Photos





User taps  
'See related  
content...'

This is a new page. Tapping 'Done' takes user right back to where he/she was



ABC 3G

02:32 PM



Done

## Related Content



# Chlamydia

### Partner Management

When patients diagnosed with chlamydia or gonorrhea indicate that their partners are unlikely to seek evaluation and treatment, providers can offer patient-delivered partner therapy (PDPT), a form of expedited partner therapy (EPT) in which partners of infected persons are treated without previous medical evaluation or prevention counseling. Because EPT might be prohibited in some states and is the topic of ongoing legislation in others (67), providers should visit [www.cdc.gov/std/ept](http://www.cdc.gov/std/ept) to obtain updated information for their individual jurisdiction.

ABC 3G

02:32 PM



## Condition Quick Pick



# Chlamydia

[See related content ...](#)

Adolescents & Adults >

Among Infants >

Among Children >



Treatments



Description



Photos

ABC 3G

02:32 PM



## Condition Quick Pick



# Chlamydia

[See related content ...](#)

### Adolescents & Adults

Diagnostic Considerations



Treatment



Follow-Up



Management of Sex Partners



Special Considerations



Treatments




Description



Photos

Example of persistent icon w/ transparency so text beneath can be seen.

If needed, would it be possible for the user to touch anywhere on the screen (except a control), and the icon disappears until user touches non-control area on screen again? And the icon could be reset to appear once user leaves this tagged topic?




If user had picked  
Special Considerations,  
which has sub-topics











User taps  
Notifications



A user can tap a notification to go to the corresponding screen where it appears. If the notification cascades from a parent topic to its children, tapping a notification from here just takes a user to the top-most screen where it applies. If a notification is for multiple parent topics, an arrow is displayed...

'New' notifications are the ones indicated in bold w/ an icon. The user hasn't run across it yet and closed it. However, if a notification is for multiple 'parent' topics and the user has not read & closed all of them, should the text will stay bold until the notification for all parent topics has been addressed???

User cannot permanently delete a notification from here. Possibly could in settings???



If user taps a notification (on the previous screen) with an arrow displayed, he/she will come to a screen like this where the user can select a topic to which the notification applies, and see it on the corresponding screen. Topics that have not yet been accessed are in bold (e.g., **Gonorrhea** and **HPV** here).

ABC 3G

02:32 PM



## Full STD Guidelines



Summary



Introduction



Methods



Clinical Prevention Guidance



Special Populations



HIV Infection: Detection,  
Counseling, and Referral



Diseases Characterized by Genital,  
Anal, or Perianal Ulcers



Management of Persons Who Have  
a History of Penicillin Allergy



Diseases Characterized by  
Urethritis and Cervicitis



Chlamydial Infections




Gonococcal Infections



Diseases Characterized by Vaginal





ABC 3G

02:32 PM



## Full STD Guidelines



### Clinical Prevention Guidance

Overview



STD/HIV Prevention Counseling



Prevention Methods




Partner Management



Reporting and Confidentiality





This is getting more granular than the Guidelines TOC

ABC 3G

02:32 PM



## Full STD Guidelines



### Pre-exposure Vaccination

Pre-exposure vaccination is one of the most effective methods for preventing transmission of some STDs. Two human papillomavirus (HPV) vaccines are available for females aged 9-26 years to prevent cervical precancer and cancer (15,16): the quadrivalent HPV vaccine (Gardasil) and the bivalent HPV vaccine (Cervarix). Gardasil also prevents genital warts. Routine vaccination of females aged 11 or 12 years is recommended with either vaccine, as is catch-up vaccination for females aged 13-26 years. Gardasil can be administered to males aged 9-26 years to prevent genital warts (17). Details regarding HPV vaccination are available at



Notification: Pre-exposure vaccines...



ABC 3G

02:32 PM



## Full STD Guidelines



### Pre-exposure Vaccination

Pre-exposure vaccination is one of the most effective methods for preventing transmission of some STDs. Two human papillomavirus (HPV) vaccines are available for females aged 9-26 years to prevent cervical precancer and cancer (15,16): the quadrivalent HPV vaccine (Gardasil) and the bivalent HPV vaccine (Cervarix). Gardasil also prevents genital warts. Routine vaccination of females aged 11 or 12 years is recommended with either vaccine, as is catch-up vaccination for females aged 13-26 years. Gardasil can be administered to males aged 9-26 years to



Notification: Pre-exposure vaccines orem ipsum dolor sit amet, consectetur quid pro quo ipsum lorem.



ABC 3G

02:32 PM



## Full STD Guidelines




### Pre-exposure Vaccination

Pre-exposure vaccination is one of the most effective methods for preventing transmission of some STDs. Two human papillomavirus (HPV) vaccines are available for females aged 9-26 years to prevent cervical precancer and cancer (15,16): the quadrivalent HPV vaccine (Gardasil) and the bivalent HPV vaccine (Cervarix). Gardasil also prevents genital warts. Routine vaccination of females aged 11 or 12 years is recommended with either vaccine, as is catch-up vaccination for females aged 13-26 years. Gardasil can be administered to males aged 9-26 years to prevent genital warts (17). Details regarding HPV vaccination are available at



[www.cdc.gov/std/hpv](http://www.cdc.gov/std/hpv). Hepatitis B



Launch the full PDF  
(access this from  
Main Menu).  
PDF Metadata  
(size, last updated,  
etc.) could go here  
too.

