



Button at top left quickly takes user to main menu



Scrolling alphabet indexer on the right. Tap a letter on the right, and list on left scrolls to that letter (like iPhone Contacts)





Contact info and
EULA could go under
About Us

Swipe the partially
hidden screen
to return to it (or
tap it on menu)

2 new
notifications

If we add the Bookmarks and Search options, they will probably go to the top





Look up a condition:

B

Bacterial Vaginosis

C

Candidiasis, Vulvovaginal

Cervicitis

Chancroid

Chlamydia

E

Epididymitis

G

Genital Warts

A
B
C
D
E
F
G
H
I
J
K
L
M
N
O
P
Q
R
S
T
U
V
W
X
Y
Z

User selects Chlamydia



Modal notification popover with corresponding 'lit' icon. Once user taps the close button, it disappears and icon becomes 'unlit'.



'Unlit' icon appears instead of lit one. Tapping this makes notification appear on the screen again.

ABC 3G

02:34 PM



Condition Quick Pick



Chlamydia



[See related content ...](#)

Adolescents & Adults

Recommended Regimens

(text)

Alternative Regimens

(text)




Treatments



Description




Photos



User taps
'See related
content...'

This is a new page. Tapping 'Done' takes user right back to where he/she was



ABC 3G

02:34 PM



Done

Related Content



Chlamydia

Partner Management

When patients diagnosed with chlamydia or gonorrhea indicate that their partners are unlikely to seek evaluation and treatment, providers can offer patient-delivered partner therapy (PDPT), a form of expedited partner therapy (EPT) in which partners of infected persons are treated without previous medical evaluation or prevention counseling. Because EPT might be prohibited in some states and is the topic of ongoing legislation in others (67), providers should visit www.cdc.gov/std/ept to obtain updated information for their individual jurisdiction.

ABC 3G

02:34 PM



Condition Quick Pick



Chlamydia



[See related content ...](#)

Adolescents & Adults >

Among Infants >

Among Children >



Treatments



Description



Photos

ABC 3G

02:34 PM



Condition Quick Pick



Chlamydia



[See related content ...](#)

Adolescents & Adults

Diagnostic Considerations



Treatment



Follow-Up



Management of Sex Partners



Special Considerations



Treatments



Description



Photos

ABC 3G

02:34 PM



Condition Quick Pick



Chlamydia



[See related content ...](#)

Diagnostic Considerations

C. trachomatis urogenital infection in women can be diagnosed by testing urine or by collecting swab specimens from the endocervix or vagina. Diagnosis of *C. trachomatis* urethral infection in men can be made by testing a urethral swab or urine specimen. Rectal *C. trachomatis* infections in persons that engage in receptive anal intercourse can be diagnosed by testing a rectal swab specimen. NAATs, cell culture, direct immunofluorescence, EIA, and nucleic acid hybridization tests are available for the



Treatments



Description



Photos

If user had picked
Special Considerations,
which has sub-topics





ABC 3G

02:34 PM



Condition Quick Pick



Chlamydia

(Chlamydia photo(s) here)




Treatments



Description



Photos




User taps
Notifications



'New' notifications are the ones indicated in bold w/ an icon. The user hasn't run across it yet and closed it. However, if a notification is for multiple 'parent' topics and the user has not read & closed all of them, should the text will stay bold until the notification for all parent topics has been addressed???

A user can tap a notification to go to the corresponding screen where it appears. If the notification cascades from a parent topic to its children, tapping a notification from here just takes a user to the top-most screen where it applies. If a notification is for multiple parent topics, an arrow is displayed...

User cannot permanently delete a notification from here. Possibly could in settings???



If user taps a notification (on the previous screen) with an arrow displayed, he/she will come to a screen like this where the user can select a topic to which the notification applies, and see it on the corresponding screen. Topics that have not yet been accessed are in bold (e.g., **Gonorrhea** and **HPV** here).

ABC 3G


02:34 PM



Full STD Guidelines



Summary	>
Introduction	>
Methods	>
Clinical Prevention Guidance	>
Special Populations	>
HIV Infection: Detection, Counseling, and Referral	>
Diseases Characterized by Genital, Anal, or Perianal Ulcers	>
Management of Persons Who Have a History of Penicillin Allergy	>
Diseases Characterized by Urethritis and Cervicitis	>
Chlamydial Infections	>
Gonococcal Infections	>



ABC 3G

02:34 PM



Full STD Guidelines



Clinical Prevention Guidance

Overview



STD/HIV Prevention Counseling



Prevention Methods




Partner Management



Reporting and Confidentiality





ABC 3G

02:35 PM



Full STD Guidelines



Prevention Methods

Abstinence & Reduction of number
of Sex Partners >

Pre-exposure Vaccination >

Male Condoms >

Female Condoms >

Cervical Diaphragms >

Topical Microbicides & Spermicides >

Condoms & n-9 Vaginal Spermicides >

Rectal Use of n-9 Spermicides >

Nonbarrier Contraception, Surgical
Sterilization, and Hysterectomy >

Male Circumcision >

Emergency Contraception (EC) >

This is getting more
granular than the
Guidelines TOC

ABC 3G

02:35 PM



Full STD Guidelines



Pre-exposure Vaccination



See related content ...

Pre-exposure vaccination is one of the most effective methods for preventing



Notification: Pre-exposure vaccines
orem ipsum dolor sit amet,
consectetur quid pro quo ipsum lorem.



To view again later, tap



Routine vaccination of females aged 11 or 12 years is recommended with either vaccine, as is catch-up vaccination for females aged 13-26 years. Gardasil can be administered to males aged 9-26 years to prevent genital warts (17). Details regarding HPV vaccination are available at www.cdc.gov/std/hpv. Hepatitis B

ABC 3G

02:35 PM



Full STD Guidelines




Pre-exposure Vaccination



[See related content ...](#)

Pre-exposure vaccination is one of the most effective methods for preventing transmission of some STDs. Two human papillomavirus (HPV) vaccines are available for females aged 9-26 years to prevent cervical precancer and cancer (15,16): the quadrivalent HPV vaccine (Gardasil) and the bivalent HPV vaccine (Cervarix). Gardasil also prevents genital warts. Routine vaccination of females aged 11 or 12 years is recommended with either vaccine, as is catch-up vaccination for females aged 13-26 years. Gardasil can be administered to males aged 9-26 years to prevent genital warts (17). Details regarding HPV vaccination are available at www.cdc.gov/std/hpv. Hepatitis B



Launch the full PDF
(access this from
Main Menu).
PDF Metadata
(size, last updated,
etc.) could go here
too.

